

INTEGRABLE $\widehat{\mathfrak{sl}_2}$ -MODULES AS INFINITE TENSOR PRODUCTS

B.FEIGIN AND E.FEIGIN

ABSTRACT. In this paper we develop an ideas and methods from [1]. Using the fusion product of the representations of the Lie algebra $\mathfrak{sl}_2$, we construct a set of the integrable highest weight $\widehat{\mathfrak{sl}_2}$ -modules L^D , depending on the vector $D \in \mathbb{N}^{k+1}$. In a special cases of D our modules are isomorphic to the irreducible $\widehat{\mathfrak{sl}_2}$ -modules $L_{i,k}$. We construct a basis of the L^D and study the decomposition of L^D on the irreducible components. We also write down a formulas for the characters of L^D .

1. INTRODUCTION

In this paper we introduce a way to construct $\widehat{\mathfrak{sl}_2}$ -modules as a limit of the finite-dimensional representations of the polynomial current Lie algebra. We use the notion of the fusion product of the finite-dimensional representations of $\mathfrak{sl}_2$, introduced in [2] and studied in [1]. Let us recall main definitions.

Let $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ be the polynomial current algebra, e, f, h – a standard $\mathfrak{sl}_2$ -basis. Let π be an $\mathfrak{sl}_2$ -module, $z \in \mathbb{C}$. Define an $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -action on the π : $(x \otimes t^i) \cdot v = z^i x \cdot v$, $x \in \mathfrak{sl}_2, v \in \pi$. Denote this $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -module by $\pi(z)$.

Let $\pi_1, \dots, \pi_n$ be an irreducible representations of $\mathfrak{sl}_2$, $\dim \pi_i = a_i$. Let $Z = (z_1, \dots, z_n) \in \mathbb{C}^n$, all z_i are pairwise distinct. Consider the tensor product of the $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -modules $\pi_1(z_1) \otimes \dots \otimes \pi_n(z_n)$. If w is a product of the highest vectors of π_i , then $\pi_1(z_1) \otimes \dots \otimes \pi_n(z_n)$ is a cyclic $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -module with a cyclic vector w ,

$$\pi_1(z_1) \otimes \dots \otimes \pi_n(z_n) = U(\mathfrak{sl}_2 \otimes \mathbb{C}[t])w.$$

Denote by $\pi_1 * \dots * \pi_n$ an adjoint graded module with respect to the filtration by the t -degree on $U(\mathfrak{sl}_2 \otimes \mathbb{C}[t])$. We call this module the fusion product of $\pi_1, \dots, \pi_n$.

In the paper we prove the existence of injective homomorphisms between different fusion products. Let $a_1 \leq \dots \leq a_n$. Then, if $a_i = a_{i+1}$, we construct an injection

$$(1) \quad \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_{i-1}} * \mathbb{C}^{a_{i+2}} * \dots * \mathbb{C}^{a_n} \hookrightarrow \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n}.$$

Moreover, there exists an isomorphism of the $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -modules

$$(2) \quad (\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n}) / (\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_{i-1}} * \mathbb{C}^{a_{i+2}} * \dots * \mathbb{C}^{a_n}) \cong \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_{i-1}} * \mathbb{C}^{a_i-1} * \mathbb{C}^{a_{i+1}+1} * \mathbb{C}^{a_{i+2}} * \dots * \mathbb{C}^{a_n}.$$

(Note that the similar formulas were used by Shilling and Warnaar to construct the supernomials, see [4, 5, 11]). Now, suppose $a_1 < a_2$. Then we construct an injection

$$(3) \quad \mathbb{C}^{a_2-a_1+1} * \mathbb{C}^{a_3} * \dots * \mathbb{C}^{a_n} \hookrightarrow \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n}$$

with a quotient:

$$(4) \quad (\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n}) / (\mathbb{C}^{a_2-a_1+1} * \mathbb{C}^{a_3} * \dots * \mathbb{C}^{a_n}) \cong \mathbb{C}^{a_1-1} * \mathbb{C}^{a_2+1} * \mathbb{C}^{a_3} * \dots * \mathbb{C}^{a_n}.$$

In fact, the more general statement is true. Let $i < j$. Then there exist submodules $S_{i,j}(A) = S_{i,j}(a_1, \dots, a_n)$ of the fusion product of $\mathbb{C}^{a_i}$, such that the following holds:

$$(5) \quad \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} / S_{i,j}(A) \cong \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_{i-1}} * \mathbb{C}^{a_i-1} * \mathbb{C}^{a_{i+1}} * \dots * \mathbb{C}^{a_{j-1}} * \mathbb{C}^{a_j+1} * \mathbb{C}^{a_{j+1}} * \dots * \mathbb{C}^{a_n}.$$

Looking at the formula (4), one can expect $S_{i,j}(A)$ to be equal to the following fusion product:

$$(6) \quad S_{i,j}(A) \cong \mathbb{C}^{a_j-a_i+1} * \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_{i-1}} * \mathbb{C}^{a_{i+1}} * \dots * \mathbb{C}^{a_{j-1}} * \mathbb{C}^{a_{j+1}} * \dots * \mathbb{C}^{a_n}.$$

But that is not true in the case of the general i, j . However, the structure of the $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -modules $S_{i,j}(A)$ is very interesting. We do not study it in this paper, but we hope to return to this topic later.

One can consider the formulas (1),(2) and (3),(4) as an equations for the tensor product of irreducible $\mathfrak{sl}_2$ -modules. To be specific,

$$(7) \quad \mathbb{C}^a \otimes \mathbb{C}^a = \mathbb{C} \oplus \mathbb{C}^{a-1} \otimes \mathbb{C}^{a+1} \text{ for } (1), (2);$$

$$(8) \quad \mathbb{C}^a \otimes \mathbb{C}^b = \mathbb{C}^{b-a+1} \oplus \mathbb{C}^{a-1} \otimes \mathbb{C}^{b+1}, a < b \text{ for } (3), (4).$$

The important special case of (1) is the following injection:

$$(9) \quad \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \hookrightarrow \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} * \mathbb{C}^{a_n} * \mathbb{C}^{a_n}.$$

Using (9), we define an inductive limit

$$(10) \quad \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} * (\mathbb{C}^{a_n})^{*2\infty} = \lim_{s \rightarrow \infty} \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} * (\mathbb{C}^{a_n})^{*2s}.$$

By definition, $\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} * (\mathbb{C}^{a_n})^{*2\infty}$ is $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -module. We will show (see proposition 3.2) that in fact, it has a structure of the representation of $\widehat{\mathfrak{sl}_2} = \mathfrak{sl}_2 \otimes \mathbb{C}[t, t^{-1}] \oplus c\mathbb{C}$, where c – a central element – acts by the multiplication on $a_n - 1$. In order to see the connection between $\widehat{\mathfrak{sl}_2}$ -modules and the fusion product, let us describe the latter in terms of the irreducible representations of $\widehat{\mathfrak{sl}_2}$.

Recall, the Lie algebra $\widehat{\mathfrak{sl}_2} = \mathfrak{sl}_2 \otimes \mathbb{C}[t, t^{-1}] \oplus c\mathbb{C}$, with the following commutator (here $(\cdot, \cdot)$ is a Killing form on $\mathfrak{sl}_2$, $x_i = x \otimes t^i$):

$$[c, x_i] = 0, [x_i, y_j] = [x, y]_{i+j} + c \cdot (x, y) i \delta_{i+j, 0}, \quad x, y \in \mathfrak{sl}_2.$$

Let $L_{i,k}$, $i = 0, \dots, k$ be an irreducible highest weight $\widehat{\mathfrak{sl}_2}$ -module with a highest vector $v_{i,k}$ such that $L_{i,k} = U(\widehat{\mathfrak{sl}_2})v_{i,k}$ and

$$(11) \quad e_{\leq 0}v_{i,k} = f_{< 0}v_{i,k} = h_{< 0}v_{i,k} = 0, \quad cv_{i,k} = kv_{i,k}, h_0v_{i,k} = iv_{i,k}.$$

In addition, there exists an operator d acting on $L_{i,k}$, defined by $[d, x_i] = ix_i$, $dv_{i,k} = 0$. It is clear that $L_{i,k}$ is bigraded by the action of the operators h_0, d . Let us denote the h_0 -grading by $\deg_z$ and d -grading by $\deg_d$.

Recall that there exists a set of the extremal vectors $v(2i)$, $i \in \mathbb{Z}$ in $L_{0,1}$, such that

$$v(0) = v_{0,1}, \quad e_{2i+1}v(2i) = v(2i+2), \quad f_{-2i+1}v(2i) = v(2i-2).$$

$L_{1,1}$ also contains a set of the extremal vectors $v(2i+1)$, $i \in \mathbb{Z}$:

$$v(1) = v_{1,1}, \quad e_{2i}v(2i-1) = v(2i+1), \quad f_{-2i}v(2i+1) = v(2i-1).$$

For the positive integers $b_1 \geq \dots \geq b_k$ define a subspace $M(b_1, \dots, b_k)$ of $L_{p_1,1} \otimes \dots \otimes L_{p_k,1}$, where p_i are 0 or 1, and parities of p_i and b_i coincide. Namely,

$$M(b_1, \dots, b_k) = U(\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}]) (v(-b_1) \otimes \dots \otimes v(-b_k)).$$

It turns out that $M(b_1, \dots, b_k)$ is isomorphic to some fusion product. To make a precise statement, let us note that by definition, a fusion product is $\mathfrak{sl}_2 \otimes \mathbb{C}[t]$ -module. But in what follows we will consider our fusions as $\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}]$ -modules (replacing $t \rightarrow t^{-1}$). Taking this remark into account, we can make the following statement:

Statement 1.1.

$M(b_1, \dots, b_k) \cong (\mathbb{C}^2)^{*b_1-b_2} * (\mathbb{C}^3)^{*b_2-b_3} * \dots * (\mathbb{C}^k)^{*b_{k-1}-b_k} * (\mathbb{C}^{k+1})^{*b_k}$
as $\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}]$ -modules.

For $A = (a_1, \dots, a_n)$ introduce an associated set

$$D = (d_1, \dots, d_{k+1}) \in \mathbb{N}^{k+1} : d_i = \#\{j : a_j = i\}, k+1 = a_n = \max\{a_i\}.$$

Denote the limit from (10) as L^D :

$$(12) \quad L^D = \lim_{s \rightarrow \infty} (\mathbb{C}^1)^{*d_1} * (\mathbb{C}^2)^{*d_2} * \dots * (\mathbb{C}^{a_n-1})^{*d_{a_n-1}} * (\mathbb{C}^{a_n})^{*(d_{a_n}+2s)}$$

(we can consider d_{a_n} to be equal to 0 or 1). We will show that L^D is a submodule in the tensor product of the irreducible level 1 $\mathfrak{sl}_2$ -modules, generated by the $\mathfrak{sl}_2$ action from some special vector. In order to prove it, we will use the realization of the irreducible level 1 representations in the fermionic space (in the space of the semi-infinite forms). Let us give a brief description of the latter.

Let $\psi(i), \phi(j), i, j \in \mathbb{Z}$ be an anticommuting variables (fermions). In particular, $\psi(i)^2 = \phi(j)^2 = 0$. The basis of F form so-called "semi-infinite" monomials in the variables $\phi(i), \psi(j)$. They are the expressions of the form

$$(13) \quad \dots \psi(N-1)\phi(N-1)\psi(N)\phi(N)\psi(i_1)\dots\psi(i_k)\phi(j_1)\dots\phi(j_l), \\ N < i_1 < \dots < i_k, N < j_1 < \dots < j_l.$$

Operators $\psi(i), \phi(j)$ acts on F by the multiplication on the corresponding fermion. Also there are the differential operators $\psi(i)^*, \phi(j)^*$, acting on F by the differentiation with respect to the variables $\psi(-j), \phi(-i)$ respectively. Thus, the following commutation relations are true ($[a, b]_+ = ab + ba$):

$$[\psi^*(i), \psi(j)]_+ = \delta_{i+j,0}, [\phi^*(i), \phi(j)]_+ = \delta_{i+j,0}.$$

The $\widehat{\mathfrak{sl}_2}$ -action on F is the following (we use the notation $a(z) = \sum_{i=-\infty}^{\infty} a_i z^i$):

$$e(z) = \psi(z)\phi(z), f(z) = \psi^*(z)\phi^*(z), h(z) = : \phi(z)\phi^*(z) : + : \psi(z)\psi^*(z) :, c = \text{Id}.$$

Define a set of vectors $v(i) \in F$:

$$v(2N) = \dots \psi(N-1)\phi(N-1)\psi(N); \quad v(2N+1) = \dots \psi(N-1)\phi(N-1)\psi(N)\phi(N).$$

We will show that there is an embedding $\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \hookrightarrow F^{\otimes(a_n-1)}$. To be precise,

$$(14) \quad \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \cong U(\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}]) (v(-d_2 - \dots - d_{a_n}) \otimes \dots \otimes v(-d_{a_n}))$$

(see statement 1.1).

Using this fermionic realization, one can see that for the irreducible $\widehat{\mathfrak{sl}}_2$ -modules we have (see also [2]):

$$L_{i,k} = \mathbb{C}^{i+1} * (\mathbb{C}^{k+1})^{*2\infty}$$

as a subspaces of $F^{\otimes k}$. Really, it is easy to see that $L_{i,k}$ can be embedded in $F^{\otimes k}$ as a union of the "pieces" of the form (14). Actually, for any D the module L^D is a subspace of $F^{\otimes k}$.

Thus, having a vector $D \in \mathbb{N}^{k+1}$, we construct the level k $\widehat{\mathfrak{sl}}_2$ -module L^D . In our paper we find a basis of L^D . Let us explain the construction of the basis in the case of the vacuum irreducible representation $L_{0,k}$.

Let $v_{0,k}$ be the highest vector in $L_{0,k}$. We realize $L_{0,k}$ as a union of the cyclic $\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}]$ -modules with a cyclic vectors $w_s, s = 1, 2, \dots$. In our special case

$$w_s = f_1^k f_3^k \dots f_{2s-1}^k v_{0,k}.$$

Denote by $e^i(j)$ the coefficient in front of z^j in the series $e(z)^i$. We prove that the union ($s = 1, 2, \dots$) of the following vectors form the $L_{0,k}$ basis:

$$(15) \quad \begin{aligned} e^k(I^k) \dots e(I^1) w_s &= e^k(i_1^k) \dots e^k(i_{l_k}^k) e^{k-1}(i_1^{k-1}) \dots e^{k-1}(i_{l_{k-1}}^{k-1}) \dots e(i_1^1) \dots e(i_{l_1}^1) w_s, \\ i_j^\alpha + 2\alpha &\leq i_{j+1}^\alpha, \\ i_j^\alpha &\geq (-2s+1)\alpha + \alpha(l_k + \dots + l_{\alpha+1})(l_k + \dots + l_{\alpha+1} + 1). \end{aligned}$$

As any integrable highest weight $\widehat{\mathfrak{sl}}_2$ -module, L^D can be decomposed into the sum of the irreducible components. Thus, one has the decomposition:

$$L^D = M_1 \otimes L_{0,k} \oplus \dots \oplus M_{k+1} \otimes L_{k,k}, \quad \dim M_i = c_{i,D}.$$

We show that M_i are graded spaces and numbers $c_{i,D}$ can be found in terms of the Verlinde algebra V_{k+1} , associated with $\mathfrak{sl}_2$. In addition, it follows from our results that the characters of M_i are the restricted Kostka polynomials (see [10], [11]). Let us recall the definition of the Verlinde algebra.

Consider an algebra with a basis $\pi_1, \pi_2, \dots$ and multiplication ($i \leq j$) $\pi_i \pi_j = \pi_{j-i+1} + \dots + \pi_{i+j-1}$ (our generators multiplies as a finite-dimensional irreducible $\mathfrak{sl}_2$ -modules). By definition, $V_k = \langle \pi_1, \pi_2, \dots \rangle / (\pi_{k+1})$. We show that in V_{k+1} the following equation is true:

$$\pi_D = \pi_1^{d_1} \dots \pi_k^{d_k} \pi_{k+1}^{d_{k+1}} = \sum_{i=1}^{k+1} c_{i,D} \pi_i.$$

The proof consists of the checking that the defining relations of the Verlinde algebra hold for our modules L^D . To be specific, we check that

$$(16) \quad \text{if } \pi_D = \pi_{D'} + \pi_{D''}, \text{ then } L^D = L^{D'} \oplus L^{D''}$$

for some choice of D, D', D'' . The set of the triples D, D', D'' is suggested by the formulas (2), (4).

Using the character formula from [1], we obtain the formulas for the character of $\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n}$ and L^D . In particular, this gives us the formula for the characters of the irreducible representations $L_{i,k}$ from [6]. Using the characters of L^D , we obtain some part of the equations $L^D = L^{D'} \oplus L^{D''}$ by the only combinatorial methods.

In the end, let us recall the method of studying of fusion products from [1]. We use the below construction as one of the main methods.

Recall that

$$\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \cong U(\mathfrak{sl}_2 \otimes \mathbb{C}[t^{-1}])w.$$

One can see that all module is generated by the only action of the operators $e_0, \dots, e_{-n+1}$. Thus, we have

$$\mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \cong \mathbb{C}[e_{-n+1}, \dots, e_0]/I^A = M^A,$$

where I^A is some ideal in the ring $\mathbb{C}[e_{-n+1}, \dots, e_0]$. Consider the rings isomorphism:

$$opp : \mathbb{C}[e_{-n+1}, \dots, e_0] \rightarrow \mathbb{C}[e_0, \dots, e_{n-1}], \quad opp(e_i) = e_{i+n-1}.$$

Denote

$$J^A = opp(I^A), W^A = \mathbb{C}[e_0, \dots, e_{n-1}]/J^A.$$

Then the following defining relations are true in W^A (here $e^{(n)}(z) = \sum_{i=0}^{n-1} e_i z^i$):

$$e^{(n)}(z)^i \div z^{\sum_{j=1}^n (i+1-a_j)_+}, \quad i = 1, 2, \dots,$$

where $s_+ = 0$ if $s \leq 0$; $s_+ = s$ if $s > 0$; and for polynomials p, q we write $p \div q$ if p is divisible on q . Our condition means that the first $-1 + \sum_{j=1}^n (i+1-a_j)_+$ coefficients of $e^{(n)}(z)^i, i = 1, 2, \dots$, generate the ideal J^A .

Our work is organized in the following way:

In the second section we construct the fermionic space F as $\widehat{\mathfrak{sl}}_2$ -module (subsection 2.1) and describe the inclusion of W^A in $F^{\otimes k}$ (theorem 2.1).

In the third section we construct $\widehat{\mathfrak{sl}}_2$ -module L^D (proposition 3.1) and describe its basis (theorem 3.1).

In the fourth section we prove the decomposition formula $L^D = L^{D'} \oplus L^{D''}$ (theorem 4.1), and also establish the connection between our modules and Verlinde algebra (proposition 4.2).

In the last section we write the character formula for L^D (theorem 5.1) and as a corollary we obtain a formula for the characters of the irreducible representations $L_{i,k}$ (corollary 5.1). In addition, we prove by the combinatorial means the equation $L^D = L^{D'} \oplus L^{D''}$ (proposition 5.1).

Remark 1.1. *In the terminology of the works [4],[5] the theorem 5.1 contains a "fermionic" formula for the character of L^D . There also exists a "bosonic" (alternating) formula, which is connected with a geometry of the flag manifolds. We hope to return to this topic later.*

Acknowledgements. The first author was partially supported by the following grants: CRDF RP1-2254, INTAS 00-55, RFBR 00-15-96579.

2. FERMIONIC PICTURE

2.1. Fermionic space (the space of the semi-infinite forms).

Consider two set of variables $\psi(i), \phi(j)$, $i, j \in \mathbb{Z}$, such that for all $i, j \in \mathbb{Z}$ we have

$$(17) \quad \begin{aligned} \psi(i)\psi(j) &= -\psi(j)\psi(i), \quad \phi(i)\phi(j) = -\phi(j)\phi(i), \\ \psi(i)\phi(j) &= -\phi(j)\psi(i), \quad \psi(i)\psi(i) = \phi(i)\phi(i) = 0. \end{aligned}$$

Let F be a linear space with a basis

$$(18) \quad \begin{aligned} &\dots \psi(N-2)\phi(N-2)\psi(N-1)\phi(N-1)\psi(N)\phi(N)\psi(i_1)\dots\psi(i_s)\phi(j_1)\dots\phi(j_t), \\ &s, t \in \mathbb{N}, \quad N, i_\alpha, j_\beta \in \mathbb{Z}, \quad N < i_1 < \dots < i_s, \quad N < j_1 < \dots < j_t. \end{aligned}$$

In other words, we consider the space, spanned by such infinite monomials in the variables $\phi(i), \psi(i)$ that there exists an integer N , such that all the $\phi(\leq N), \psi(\leq N)$ are factors in it. In addition, numbers of all factors are less than some natural number. Let us call the monomials with the above properties configurations, and let the product $\prod_{i=-\infty}^N \psi(i)\phi(i)$ be a tale of the configuration. It is clear that the fermions, which are not in the tale, can be reordered, using the commutation relations.

Define an action of the operators $\psi(i), \phi(i), \psi^*(i), \phi^*(i)$ on F . By the definition, $\psi(i)(v)$ ($\phi(i)(v)$), where v is a configuration, is a monomial $v\psi(i)$ ($v\phi(i)$) (to get a basis element probably one must reorder the fermions). For example, if v contains $\psi(i)$ ($\phi(i)$), then $\psi(i)v = 0$ ($\phi(i)v = 0$).

In its turn, the operators $\psi^*(i), \phi^*(i)$ act by the differentiation with respect to the variables $\psi(-i), \phi(-i)$ respectively. The following commutation relations hold ($[a, b]_+ = ab + ba$):

$$(19) \quad [\psi^*(i), \psi(j)]_+ = \delta_{i+j,0}, \quad [\phi^*(i), \phi(j)]_+ = \delta_{i+j,0}, \\ [\psi^*(i), \phi(j)] = [\phi^*(i), \psi(j)] = 0.$$

In addition, all $\psi^*(i), \phi^*(j)$ mutually anticommute.

Introduce the generating functions:

$$(20) \quad \psi(z) = \sum_{i=-\infty}^{\infty} \psi(i)z^i, \quad \phi(z) = \sum_{i=-\infty}^{\infty} \phi(i)z^i, \\ \psi^*(z) = \sum_{i=-\infty}^{\infty} \psi^*(i)z^i, \quad \phi^*(z) = \sum_{i=-\infty}^{\infty} \phi^*(i)z^i.$$

Recall the construction of the $\widehat{\mathfrak{sl}}_2 = \mathfrak{sl}_2 \otimes \mathbb{C}[z, z^{-1}] + c\mathbb{C}$ action on F . The central element acts by 1 (F is a level 1 representation). Define an action of the generating functions of the elements e, f, h on F as follows (for $x \in \mathfrak{sl}_2$ denote $x_i = x \otimes z^i$):

$$(21) \quad e(z) = \sum_{i=-\infty}^{\infty} e_i z^i = \psi(z)\phi(z); \quad f(z) = \sum_{i=-\infty}^{\infty} f_i z^i = \psi^*(z)\phi^*(z),$$

and the action of $h(z)$ can be defined using the commutation relations in $\widehat{\mathfrak{sl}}_2$.

Introduce a set of vectors $v(i) \in F$:

$$(22) \quad v(2N) = \dots \psi(N-2)\phi(N-2)\psi(N-1)\phi(N-1)\psi(N); \\ v(2N+1) = \dots \psi(N-1)\phi(N-1)\psi(N)\phi(N).$$

Note that $e_{N+1}v(N) = v(N+2), e_{\leq N}v(N) = 0$.

Note also that the space $\underbrace{F \otimes \dots \otimes F}_k$ inherit a structure of level k $\widehat{\mathfrak{sl}}_2$ -module.

2.2. A fermionic realization of $\mathbb{C}[e_0, \dots, e_{n-1}]$ -modules.

Recall that in [1] we define a set of $\mathbb{C}[e_0, \dots, e_{n-1}]$ -modules W^A , depending on the vector $A \in \mathbb{N}^n$: $W^A = \mathbb{C}[e_0, \dots, e_{n-1}]/J^A$, where J^A is an ideal in the ring $\mathbb{C}[e_0, \dots, e_{n-1}]$, generated by the conditions on the generating function $e^{(n)}(z) = \sum_{i=0}^{n-1} e_i z^i$:

$$(23) \quad e^{(n)}(z)^i \div z^{\sum_{j=1}^n (i+1-a_j)_+}$$

(we use the notation $s_+ = 0$, if $s < 0$, and $s_+ = s$, if $s \geq 0$; $p \div q$ means that p is divisible on q). In other words, the ideal J^A is generated by the first $-1 + \sum_{j=1}^n (i + 1 - a_j)_+$ coefficients of the polynomials $e^{(n)}(z)^i, i = 1, 2, \dots$

Let $A = (a_1, \dots, a_n)$ and $a_1 \leq \dots \leq a_n$. In [1] the following statement was proved:

Statement 2.1. $W^{a_1, \dots, a_{n-1}} \hookrightarrow W^A$ and $W^A / W^{a_1, \dots, a_{n-1}} \cong W^{a_1, \dots, a_{n-1}, a_n - 1}$.

By induction, one can obtain that $\dim W^A = \prod_{i=1}^n a_i$.

Let $k+1 = a_n = \max\{a_i : i = 1, \dots, n\}$, $d_j = \#\{i : a_i = j\}, j = 1, \dots, k+1$. We say that the set $D = (d_1, \dots, d_{k+1})$ is associated with A . Define a vector $v_A \in F^{\otimes k}$:

$$(24) \quad v_A = v(-1 + d_1) \otimes v(-1 + d_1 + d_2) \otimes \dots \otimes v(-1 + d_1 + \dots + d_k).$$

Denote $\widetilde{W}^A = \mathbb{C}[e_0, \dots, e_{n-1}](v_A)$, $\widetilde{W}^A \hookrightarrow F^{\otimes k}$. Let us prove the following inclusion theorem:

Theorem 2.1. $\widetilde{W}^A \cong W^A$ as $\mathbb{C}[e_0, \dots, e_{n-1}]$ -modules.

Proof. First, we prove that in $F^{\otimes k}$

$$e^{(n)}(z)^i(v_A) \div z^{\sum_{j=1}^n (i+1-a_j)_+}$$

(in other words, first $\sum_{j=1}^n (i+1-a_j)_+ - 1$ coefficients of $e^{(n)}(z)^i(v_A)$ are equal to zero). Note that

$$e(z)^i v_A \div z^{\sum_{j=1}^n (i+1-a_j)_+}, \text{ if } i \leq k; \quad e(z)^i v_A = 0 \text{ if } i > k$$

(first is a consequence of the condition $e_{\leq N} v(N) = 0$, and second is true, because $e(z)^{k+1} = 0$ in $F^{\otimes k}$). So, for all i we obtain

$$e(z)^i v_A \div z^{\sum_{j=1}^n (i+1-a_j)_+}.$$

We have $(x = -\sum_{i=n}^{\infty} z^{i-n} e_i)$:

$$e^{(n)}(z)^i v_A = (e(z) + z^n x)^i v_A = \sum_{j=0}^i \binom{i}{j} (e(z)^{i-j} z^{nj} x^j v_A).$$

Note that

$$(e(z)^{i-j} z^{nj}) v_A \div z^{\sum_{l=1}^n (i+1-a_l)_+},$$

because

$$nj + \sum_{l=1}^n (i-j+1-a_l)_+ \geq \sum_{l=1}^n (i+1-a_l)_+.$$

Thus, we obtain $J^A v_A = 0$ and as a corollary, $\dim \mathbb{C}[e_0, \dots, e_{n-1}] v_A \leq \prod_{i=1}^n a_i$.

To prove our theorem, it is enough to show that $\dim \widetilde{W}^A = \prod a_i$.

In the following lemma we use the notations:

$$Aa_{n+1} = (a_1, \dots, a_n, a_{n+1}), \quad A(a_{n+1} - 1) = (a_1, \dots, a_n, a_{n+1} - 1).$$

Lemma 2.1. *Let $A = (a_1, \dots, a_n)$, $a_1 \leq \dots \leq a_n$. Let $a_{n+1} \geq a_n$. Denote $B_1 \subset \mathbb{C}[e_0, \dots, e_{n-1}]$, $B_2 \subset \mathbb{C}[e_0, \dots, e_n]$ such sets of monomials that $B_1 v_A = \{x v_A, x \in B_1\}$, $B_2 v_{A(a_{n+1}-1)} = \{x v_{A(a_{n+1}-1)}, x \in B_2\}$ are linearly independent sets of vectors in $\widetilde{W}^A$ and $\widetilde{W}^{A(a_{n+1}-1)}$ respectively. Let $e_n B_2 = \{e_n x, x \in B_2\}$. Then $B_1 v_{Aa_{n+1}} \cup (e_n B_2) v_{Aa_{n+1}}$ is a linearly independent set of vectors in $\widetilde{W}^{Aa_{n+1}}$.*

Proof. Consider 3 different cases:

$$a_{n+1} > a_n + 1, \quad a_{n+1} = a_n + 1, \quad a_{n+1} = a_n.$$

1. $a_{n+1} > a_n + 1$.

Note that in this case

$$\begin{aligned} (25) \quad v_1 &= v_A = v(-1 + d_1) \otimes \dots \otimes v(-1 + d_1 + \dots + d_{a_n-1}), \\ v_2 &= v_{A(a_{n+1}-1)} = v_A \otimes v(-1 + n)^{\otimes(a_{n+1}-a_n-1)}, \\ v_3 &= v_{Aa_{n+1}} = v_{A(a_{n+1}-1)} \otimes v(-1 + n). \end{aligned}$$

Let $b_2 \in B_2$, $b_2 = e_n^i x$, $x \in \mathbb{C}[e_0, \dots, e_{n-1}]$. Recall that

$$(26) \quad e_{<n} v(-1 + n) = 0, \quad e_n v(-1 + n) = v(1 + n), \quad e_n^2 v(-1 + n) = 0.$$

But the sum of the summands in $e_n b_2 v_3$, with a property that the last factor has been changed (i.e. it equals not $v(-1 + n)$, but $v(1 + n)$), equals to

$$(27) \quad (i + 1) b_2 v_2 \otimes v(1 + n).$$

Thus, since the set of vectors $B_2 v_2$ is linearly independent, we obtain the linearly independence of $e_n B_2 v_3$. In its turn, the linearly independence of $B_1 v_3$ is obvious, because $B_1 \subset \mathbb{C}[e_0, \dots, e_{n-1}]$ and so for $b_1 \in B_1$

$$(28) \quad b_1 v_3 = b_1 v_1 \otimes v(-1 + n)^{\otimes(a_{n+1}-a_n)}.$$

Now, we must prove that $B_1 v_3$ and $e_n B_2 v_3$ are linearly independent as a whole. But that is a consequence from the formula (28) and speculations, leading to the formula (27).

2. $a_{n+1} = a_n + 1$

In this case

$$\begin{aligned} (29) \quad v_A &= v(-1 + d_1) \otimes \dots \otimes v(-1 + d_1 + \dots + d_{a_n-1}), \\ v_{A(a_{n+1}-1)} &= v_A, \\ v_{Aa_{n+1}} &= v_A \otimes v(-1 + n). \end{aligned}$$

The proof is the same as in the previous case.

3. $a_{n+1} = a_n$

In this case

$$\begin{aligned} (30) \quad v_1 &= v_A = v(-1 + d_1) \otimes \dots \otimes v(-1 + d_1 + \dots + d_{a_n-1}), \\ v_3 &= v_{Aa_{n+1}} = v_A, \\ v_2 &= v_{A(a_{n+1}-1)} = v(-1 + d_1) \otimes \dots \otimes v(-1 + d_1 + \dots + d_{a_n-2}) \otimes \\ &\quad \otimes v(-1 + d_1 + \dots + d_{a_n-2} + d_{a_n-1} + 1). \end{aligned}$$

Let us prove that $e_n B_2 v_3$ is linearly independent. Recall that all of our vectors are the elements of the tensor powers $F^{\otimes l}$. We denote the fermions, used in the construction of the s -th factor as $\psi_s(i), \phi_s(j)$.

Consider a vector $w \in F^{\otimes(a_n-1)}$, defined as follows:

$$(31) \quad \begin{aligned} w &= \phi_{a_n-1}(n - (d_1 + \dots + d_{a_n-1})/2)v_2, \text{ if } d_1 + \dots + d_{a_n-1} \text{ is even;} \\ w &= \psi_{a_n-1}(n - (d_1 + \dots + d_{a_n-1} - 1)/2)v_2, \text{ if } d_1 + \dots + d_{a_n-1} \text{ is odd.} \end{aligned}$$

Thus, $w = \chi v_2$, where χ is defined by (31). One can see that w is a summand in $e_n v_3$. Note also that

$$(32) \quad \mathbb{C}[e_0, \dots, e_n]v_2 \cong \mathbb{C}[e_0, \dots, e_n]w.$$

In fact, we multiply v_2 on the fermion with a such number that beeing added to the maximal number of the opposite fermion (ψ is opposite to ϕ and ϕ to ψ), which is a multiplier in the last factor in v_2 , it alreedy gives n . So, χ cannot appear, while acting by the polynomials in the variables $e_0, \dots, e_n$ on v_2 .

Now, let l be a linear combination of the elements of B_2 . One can see that the summands of $e_n l \cdot v_3$, which contain χ as a multiplier in the last factor, arise only while acting e_n in the last factor of the tensor power. In addition, if $b_2 \in B_2$, $b_2 = e_n^i x$, $x \in \mathbb{C}[e_0, \dots, e_{n-1}]$, then the sum of the summands in $e_n b_2 v_3$, which contain χ in the last factor, equals $(i+1)b_2 w$. Now, we obtain the linearly independence of $e_n B_2 v_3$ as a consequence of the linearly independence of $B_2 v_2$ and equality (32).

To finish the proof of the lemma, note that, firstly, $B_1 v_3$ is linearly independent since $v_3 = v_1$, and, secondly, $B_1 v_3 \cup e_n B_2 v_3$ is linearly independent since $B_1 \subset \mathbb{C}[e_0, \dots, e_{n-1}]$ and thus χ cannot appear in $b_1 v_3$, $b_1 \in B_1$. $\square$

We have proved the lemma. Now we obtain our theorem by induction on the sum of a_i (including the existense of the monomial basis in the inductive assumption). $\square$

3. $\widehat{\mathfrak{sl}_2}$ -MODULES. BASES.

3.1. Construction of the $\widehat{\mathfrak{sl}_2}$ -modules.

Let $A = (a_1 \leq \dots \leq a_n), k+1 = a_n$. Consider the image $I^A \in \mathbb{C}[e_{-n+1}, \dots, e_0]$ of the ideal J^A under the action of the ring isomorphism

$$(33) \quad \mathbb{C}[e_0, \dots, e_{n-1}] \rightarrow \mathbb{C}[e_{-n+1}, \dots, e_0], \quad e_i \mapsto e_{i-n+1}.$$

As it was said in the introduction, the corresponding quotient is isomorphic to the fusion product of the $\mathfrak{sl}_2$ -modules. To be specific,

$$\mathbb{C}[e_{-n+1}, \dots, e_0]/I^A \cong \mathbb{C}^{a_1} * \dots * \mathbb{C}^{a_n} \text{ as } \mathfrak{sl}_2 \otimes (\mathbb{C}[z^{-1}]/z^{-n})\text{-modules}$$

(actually, $\mathbb{C}[e_{-n+1}, \dots, e_0]/I^A$ has a structure of $\mathfrak{sl}_2 \otimes (\mathbb{C}[z^{-1}]/z^{-n})$ -module, see [1]). From the theorem (2.1) we obtain the following statement:

Statement 3.1. *Let*

$$(34) \quad \begin{aligned} w_A &= v(d_1 - n) \otimes v(d_1 + d_2 - n) \otimes \dots \otimes v(d_1 + \dots + d_k - n), \\ M^A &= \mathbb{C}[e_{-n+1}, \dots, e_0]w_A. \end{aligned}$$

Then as $\mathfrak{sl}_2 \otimes (\mathbb{C}[z^{-1}]/z^{-n})$ -modules $M^A \cong \mathbb{C}[e_{-n+1}, \dots, e_0]/I^A$.

Note that since $d_1 + \dots + d_{k+1} = n$, then

$$w_A = v(-d_2 - \dots - d_{k+1}) \otimes v(-d_3 - \dots - d_{k+1}) \otimes \dots \otimes v(-d_{k+1}).$$

Denote $A_s = A \underbrace{a_n, \dots, a_n}_{2s}$, $A_s \in \mathbb{N}^{n+2s}$. We will prove that $M^{A_s} \hookrightarrow M^{A_{s+1}}$ as a subspaces of $F^{\otimes k}$.

Proposition 3.1. $w_A \in U(\mathfrak{sl}_2 \otimes \mathbb{C}[z^{-1}])w_{A_1}$.

Proof. Recall that $d_j = \#\{i : a_i = j\}$. Introduce a notation:

$$\beta_i = d_{i+1} + \dots + d_{k+1}, \quad i = 1, \dots, k.$$

Then we have

$$(35) \quad \begin{aligned} w_A &= v(-\beta_1) \otimes \dots \otimes v(-\beta_k), \\ w_{A_1} &= v(-\beta_1 - 2) \otimes \dots \otimes v(-\beta_k - 2). \end{aligned}$$

Note that the vectors

$$w_A, e_1 w_A, \dots, e_1^{\sum_{i=1}^k (\beta_i + 1)} w_A$$

form an $\mathfrak{sl}_2$ -module with respect to the $\mathfrak{sl}_2 = \langle e_1, h_0 + k \cdot \text{Id}, f_{-1} \rangle$. From one hand, it can be checked by the direct calculation; from the other one can see that there exists an isomorphism

$$\mathbb{C}[e_1, e_0, \dots, e_{-n+1}]w_A \cong \mathbb{C}[e_0, \dots, e_{-n}]/I^{A_{a_n}}, \quad e_i \mapsto e_{i-1}.$$

But in any fusion product the e_0 -action on the highest vector spans the $\mathfrak{sl}_2$ -module.

Note that

$$(36) \quad e_1^{\sum_{i=1}^k (\beta_i + 1)} w_A \text{ is proportional to } v(\beta_1 + 2) \otimes \dots \otimes v(\beta_k + 2).$$

Also, one can see that

$$(37) \quad e_0^{\sum_{i=1}^k (\beta_i + 2)} w_{A_1} \text{ is proportional to } v(2 + \beta_1) \otimes \dots \otimes v(2 + \beta_k).$$

Thus, we obtain the following equality

$$(38) \quad \text{const} \cdot w_A = f_{-1}^{\sum_{i=1}^k (\beta_i + 1)} e_0^{\sum_{i=1}^k (\beta_i + 2)} w_{A_1}$$

(of course, the constant doesn't equal to zero). Proposition is proved. $\square$

Remark 3.1. From the proof we obtain that $e_1 M_A \hookrightarrow M_{A_1}$.

Remark 3.2. From the proof of the proposition one can obtain that

$$w_{A_1} \hookrightarrow U(\mathfrak{sl}_2 \otimes \mathbb{C}[z])w_A.$$

We have proved that $w_A \in U(\mathfrak{sl}_2 \otimes \mathbb{C}[z^{-1}])w_{A_1}$. Thus, we have a chain of the inclusions:

$$M^A \hookrightarrow M^{A_1} \hookrightarrow M^{A_2} \hookrightarrow \dots$$

Definition 3.1. Let $D \in (\mathbb{N} \cup 0)^{k+1}$, and $d_{k+1} = 0$ or $d_{k+1} = 1$. Let $d_1 + \dots + d_{k+1} = n$. Consider $A \in \mathbb{N}^{n+2}$, such that $d_j = \#\{i : a_i = j\}$ for $j = 1, \dots, k$ and $\#\{i : a_i = k+1\} = d_{k+1} + 2$. Then

$$L^D = L^{d_1, \dots, d_{k+1}} = \bigcup_{s=0}^{\infty} M^{A_s}.$$

Notation:

$$L^D = (\mathbb{C}^1)^{*d_1} * (\mathbb{C}^2)^{*d_2} * \dots * (\mathbb{C}^k)^{*d_k} * (\mathbb{C}^{k+1})^{*(d_{k+1} + 2\infty)}.$$

By definitions, L^D is only $\mathfrak{sl}_2 \otimes \mathbb{C}[z^{-1}]$ -module. We show that in fact it has a structure of $\widehat{\mathfrak{sl}_2}$ -module.

Proposition 3.2. L^D is closed with respect to the action of $\widehat{\mathfrak{sl}_2}$.

Proof. By the remark (3.1), L^D is closed with respect to the action of e_1 . In addition, L^D is $\mathfrak{sl}_2 \otimes \mathbb{C}[z^{-1}]$ -module. But $\widehat{\mathfrak{sl}_2}$ is generated by $\mathfrak{sl}_2 \otimes \mathbb{C}[z^{-1}]$ and e_1 . Proposition is proved. $\square$

It is clear that $L^{d_1, \dots, d_{k+1}}$ is an integrable level k representation. It is cyclic $\widehat{\mathfrak{sl}_2}$ -module and may be generated from any of the vectors w_{A_s} (see the proposition 3.1 and remark 3.2).

Corollary 3.1. *Let $0 \leq i \leq k$. Define such D that $d_j = \delta_{i+1, j}$. Then $L^D \cong L_{i, k}$. We can write it as $L_{i, k} = \mathbb{C}^{i+1} * (\mathbb{C}^{k+1})^{*2\infty}$.*

Remark 3.3. *One can see that $L^{d_1, d_2, \dots, d_{k+1}} = L^{d_2, \dots, d_{k+1}}$. That means that the number of the one-dimensional representations is not important.*

Remark 3.4. *Let $p(D)$ be a number of the odd numbers in the sequence*

$$d_{k+1}, d_{k+1} + d_k, \dots, d_{k+1} + \dots + d_2.$$

Then

$$L^D \hookrightarrow L_{1,1}^{\otimes p(D)} \otimes L_{0,1}^{\otimes (k-p(D))}.$$

3.2. Basis of L^D .

Here we construct an L^D basis. First we will construct the basis of the space $W^{A_\infty} = \mathbb{C}[\dots, e_{-1}, e_0, e_1, \dots]v_A$.

Let $k+1 = a_n = \max\{a_i\}$. Recall that $e(z) = \sum_{i=-\infty}^{\infty} e_i z^i$. Denote by $e^i(j)$ the coefficient in $e(z)^i$ in front of z^j . Note that $e_{<0}v_A = 0$, so $e(z)v_A = (\sum_{i=0}^{\infty} e_i z^i)v_A$. Let us prove the following lemma:

Lemma 3.1. *The following elements are linearly independent in W^{A_∞} :*

$$(39) \quad e(I^1) \dots e^k(I^k)v_A = e(i_1^1) \dots e(i_{l_1}^1) e^2(i_1^2) \dots e^2(i_{l_2}^2) \dots e^k(i_1^k) \dots e^k(i_{l_k}^k)v_A,$$

$$(40) \quad i_j^\alpha + 2\alpha \leq i_{j+1}^\alpha,$$

$$i_j^\alpha \geq \alpha d_1 + (\alpha - 1)d_2 + \dots + d_\alpha + \alpha(l_k + \dots + l_{\alpha+1})(l_k + \dots + l_{\alpha+1} + 1).$$

Proof. Recall that $v_A \in F^{\otimes k}$. We denote the fermions, used in the construction of the s -th factor of $F^{\otimes k}$ as $\psi_s(i), \phi_s(j)$. Denote $e_{i,s}$ the following operators in $F^{\otimes k}$:

$$e_{i,s} = \underbrace{\text{Id} \otimes \dots \otimes \text{Id}}_{s-1} \otimes e_i \otimes \text{Id} \otimes \dots \otimes \text{Id}.$$

Let $[x]$ be an integral part of x . Define an operator

$$m(i_1^\alpha, \dots, i_{l_\alpha}^\alpha, d_1, \dots, d_\alpha) = m(I^\alpha, d_1, \dots, d_\alpha) : F^{\otimes k} \rightarrow F^{\otimes k}$$

by the following way:

$$(41) \quad \prod_{j=1}^{\alpha-1} \prod_{m=1}^{l_\alpha} e_{d_1+\dots+d_j+2(m-1), j} \times \\ \times \prod_{m=1}^{l_\alpha} \psi_\alpha \left(\left[\frac{i_m^\alpha - (\alpha-1)d_1 - (\alpha-2)d_2 - \dots - d_{\alpha-1} - (\alpha-1)2(m-1) + 1}{2} \right] \right) \times \\ \times \prod_{m=1}^{l_\alpha} \phi_\alpha \left(\left[\frac{i_m^\alpha - (\alpha-1)d_1 - (\alpha-2)d_2 - \dots - d_{\alpha-1} - (\alpha-1)2(m-1)}{2} \right] \right).$$

Let $m(I^1, \dots, I^k)v_A \in F^{\otimes k}$ be the following vector:

$$(42) \quad \begin{aligned} m(I^1, \dots, I^k)v_A &= m(I^1, d_1 + 2(l_k + \dots + l_2))m(I^2, d_1 + 2(l_k + \dots + l_3), d_2) \dots \\ &\dots m(I^{k-1}, d_1 + 2l_k, d_2, \dots, d_{k-1})m(I^k, d_1, \dots, d_k)v_A. \end{aligned}$$

Note that if $I^1, \dots, I^k$ satisfy the condition (40), then $m(I^1, \dots, I^k)v_A$ is one of the summands in $e(I^1) \dots e^k(I^k)v_A$. Let us explain why.

Let us act in a following way: first by $e^k(I^k)$, then by $e^{k-1}(I^{k-1})$, and so on till to $e(I^1)$. All this operators act on $F^{\otimes k}$. Every time, while acting by $e^\alpha(I^\alpha)$ on $F^{\otimes k}$ we consider only the summands in which the action is happening on the first α factors. In addition, the action on the first $\alpha - 1$ factors is happening by the "minimal" possible way, that is we are multiplying on the fermions with a minimal possible numbers. In the same time the action on the α -th factor is happening by the "middle" way, that is $e^\alpha(j)$ acts on the α -th factor by the multiplication on $\psi_\alpha[\frac{j+1}{2}]\phi_\alpha[\frac{j}{2}]$. Condition (40) guaranties that while acting by the described way, we will get a nonzero vector.

Now, let us prove the linearly independence of our vectors. Consider the linear combination

$$(43) \quad \sum \beta_{I^1, \dots, I^k} e(I^1) \dots e^k(I^k)v_A,$$

where all the sets $(I^1, \dots, I^k)$ satisfy the condition (40). Between the sets from (43) choose the one $(I_0^1, \dots, I_0^k)$ with a property that for all other sets from (43), we have: there exists j, m , such that $(i_0)_m^j > i_m^j$ and for all such j_1, m_1 , that $j_1 > j$ or $j_1 = j, m_1 < m$ we have $(i_0)_{m_1}^{j_1} = i_{m_1}^{j_1}$. Note that from the construction, we obtain that $m(I_0^1, \dots, I_0^k)v_A$ appears in (43) only from $e(I_0^1) \dots e^k(I_0^k)v_A$ (and with a nonzero coefficient) and doesn't appear from any other summand. Thus we obtain that

$$\sum \beta_{I^1, \dots, I^k} e(I^1) \dots e^k(I^k)v_A \neq 0.$$

Lemma is proved. $\square$

Proposition 3.3. *Vectors (39) with a condition (40), form a base of W^{A_∞} .*

Proof. Since we have checked the linearly independence, it is enough to compare the characters of the probable basis and the space W^{A_∞} . One can see that the character of the probable basis is

$$(44) \quad \sum_{i_1, \dots, i_k=0}^{\infty} (zq^{-1})_{i_1}^{\sum_{t=1}^k l_{i_t}} \frac{q^{\sum_{s,t=1}^k \min(s,t) i_s i_t + \sum_{t=1}^k d_t(i_t + 2i_{t+1} + \dots + (k-l+1)i_k)}}{(i_1)_q! \dots (i_k)_q!}.$$

Thus, the character of W^{A_∞} is more or equal (in every homogeneous component) than the expression (44). We will show that in fact we have an equality.

Consider a quotient B^A of the polynomial ring

$$\mathbb{C}[b_1(0), b_1(1), \dots, b_2(0), b_2(1), \dots, b_k(0), b_k(1), \dots].$$

The defining relations in B^A are the following relations on the generating functions $b_i(z) = \sum_{j=0}^{\infty} b_i(j)z^j$ (we use the notation $b^{(s)}(z)$ for the s -th derivative of the

series):

$$(45) \quad \begin{aligned} b_i(z) &\div z^{id_1+(i-1)d_2+\dots+d_i}, \\ b_i^{(s)}(z)b_j^{(t)}(z) &= 0 \text{ if } i < j, \ s+t < 2i. \end{aligned}$$

Statement 3.2. Define $\deg_z b_i(j) = i$, $\deg_q b_i(j) = j$. Then the character of B^A is coinciding with the expression (44).

Proof. It is known, (see for example [9]) that the dual space $(B^A)^*$ can be realized as a sum of the spaces of polynomials

$$f(z_1^1, \dots, z_{i_1}^1; \dots; z_1^k, \dots, z_{i_k}^k), \ i_\alpha \geq 0,$$

symmetric with respect to all of the groups of variables $z_1^s, \dots, z_{i_s}^s$ and which can be written in a following form:

$$(46) \quad f = g \prod_{l=1}^k \prod_{i=1}^{i_l} (z_i^l)^{ld_1+(l-1)d_2+\dots+d_l} \prod_{1 \leq s \leq t \leq k} \prod_{\substack{i=1, \dots, i_s \\ j=1, \dots, i_t}} (z_i^s - z_j^t)^{2s},$$

where $g(z_1^1, \dots, z_{i_1}^1; \dots; z_1^k, \dots, z_{i_k}^k)$ is an arbitrary polynomial, symmetric with respect to all of the groups of variables. In other words, f is vanishing $2s$ times on the diagonals $z_i^s = z_j^t$, $s \leq t$.

One can see that the character of the polynomials (46) coincides with (44). $\square$

To prove the proposition, it is enough to show that $\text{ch} W^{A\infty} \leq \text{ch} B^A$ (we mean that the inequalities are true in each homogeneous component). In order to do it, we will use the construction for the filtration on the dual space $(W^{A\infty})^*$, which is explained in [9] in a more general case.

One can see that a dual space $(W^{A\infty})^*$ can be realized as a sum of the spaces of the polynomials $f(z_1, \dots, z_s)$, $s = 0, 1, 2, \dots$, which satisfy the following conditions:

$$(47) \quad f(\underbrace{z, \dots, z}_i, z_{i+1}, \dots, z_s) \div z^{id_1+(i-1)d_2+\dots+d_i}, \ i = 1, \dots, k,$$

$$f(\underbrace{z, \dots, z}_{k+1}, z_{k+2}, \dots, z_s) = 0.$$

(This realization is tantamount to the fact that $W^{A\infty}$ is a quotient of the polynomial ring $\mathbb{C}[e_0, e_1, \dots]$, with a defining relations:

$$(48) \quad e^i(z) \div z^{id_1+\dots+d_i}, \ i = 1, \dots, k, \ e^{k+1}(z) = 0.$$

In [1] we have proved it in the case $d_1 = \dots = d_k = 0$. In the general case the proof is the same.)

As it is shown in [9], there is a filtration on the space $(W^{A\infty})^*$ by the subspaces, enumerated by the Young diagrams, and the adjoint quotients are isomorphic to the spaces of polynomials (46). This gives us that the character of $W^{A\infty}$ is equal to (44). $\square$

Corollary 3.2. *Elements*

$$(49) \quad e(I^1) \dots e^k(I^k) w_A = e(i_1^1) \dots e(i_{l_1}^1) e^2(i_1^2) \dots e^k(i_{l_2}^2) \dots e^k(i_1^k) \dots e^k(i_{l_k}^k) w_A,$$

$$i_j^\alpha + 2\alpha \leq i_{j+1}^\alpha,$$

$$i_j^\alpha \geq \alpha - \alpha(d_{k+1} + \dots + d_{\alpha+1}) - (\alpha-1)d_\alpha - \dots - d_2 + (l_k + \dots + l_{\alpha+1})(l_k + \dots + l_{\alpha+1} + 1)$$

form a base of $\mathbb{C}[\dots, e_{-1}, e_0, e_1, \dots]w_A$. (Denote this set of vectors as B_A).

Lemma 3.2. $B_{A_s} \subset B_{A_{s+1}}$.

Proof. Note that in $F^{\otimes k}$ the following is true

$$e(z)^k = k! \cdot e(z) \otimes \dots \otimes e(z).$$

Thus we obtain the following equality ($\alpha_i = d_2 + \dots + d_i$):

$$(50) \quad e^k \left(\sum_{i=2}^{k+1} (-\alpha_i - 1) \right) w_{A_2} = k! \cdot w_A.$$

As a consequence, we obtain that $B_{A_s} \subset B_{A_{s+1}}$. $\square$

Theorem 3.1. $\bigcup_{s=0}^{\infty} B_{A_s}$ is a basis of L^D .

Proof. By definition

$$L^D = \bigcup_{s=0}^{\infty} \mathbb{C}[\dots, e_{-1}, e_0, e_1, \dots]w_{A_s}.$$

Using the lemma 3.2, we obtain the theorem. $\square$

Remark 3.5. Using the theorem 3.1 and formula (44) one can obtain the character of L^D (note that in [6] character of $L_{i,k}$ was obtained by this method). But in the last section we will obtain the character of L^D using a fact that L^D is a limit of finite-dimensional fusion products.

4. THE DECOMPOSITION OF L^D . VERLINDE ALGEBRA.

4.1. Functional recurrent formula.

Let $A = (a_1 \leq \dots \leq a_n) \in \mathbb{N}^n$, $k+1 = a_n$, $D \in (\mathbb{N} \cup 0)^{k+1}$ is associated with A . We will construct such A', A'' that $W^{A'} \hookrightarrow W^A$ and $W^A/W^{A'} \cong W^{A''}$.

Recall that $e^{(n)}(z) = \sum_{i=0}^{n-1} e_i z^i$. Denote by $[e^{(n)}(z)^i]_j$ the coefficient in front of z^j in $e^{(n)}(z)^i$.

Lemma 4.1. Let $d_l \neq 0$. Then

$$[e^{(n)}(z)^{l-1}]_{\sum_{j=1}^{l-1} (l-j)d_j} v_A = (l-1)! (e_{d_1} \otimes \dots \otimes e_{d_1+\dots+d_{l-1}} \otimes \text{Id} \otimes \dots \otimes \text{Id}) v_A.$$

Proof. Firstly, note that

$$(51) \quad [e^{(n)}(z)^{l-1}]_{\sum_{j=1}^{l-1} (l-j)d_j} v_A = e^{l-1} \left(\sum_{j=1}^{l-1} (l-j)d_j \right) v_A$$

(recall that $e^i(j)$ is a coefficient in front of z^j in the series $e(z)^i$). In fact, if $c = \sum_{i=n}^{\infty} e_i z^{i-n}$, then

$$(52) \quad e(z)^{l-1} v_A = e^{(n)}(z)^{l-1} v_A + \sum_{i=0}^{l-2} \binom{l-1}{i} z^{n(l-1-i)} c^{l-1-i} e^{(n-1)}(z)^i v_A.$$

Recall that

$$e^{(n)}(z)^i v_A \div z^{\sum_{j=1}^i d_j (i+1-j)}.$$

We have:

$$z^{n(l-1-i)} e^{(n-1)}(z)^i v_A \div z^{n(l-1-i) + \sum_{j=1}^i d_j(i+1-j)}.$$

In the same time, since $\sum_{j=1}^{k+1} d_j = n$ and $d_l \neq 0$, then

$$n(l-1-i) + \sum_{j=1}^i d_j(i+1-j) > \sum_{j=1}^{l-1} (l-j)d_j.$$

From this and (52), we obtain (51), because all the summands in the right hand side of (52), but $e^{(n)}(z)^{l-1}$ are divisable on $z^{1+\sum_{j=1}^{l-1} d_j(l-j)}$.

Consider now $e(z)^{l-1} v_A$. Since $e(z)^2 = 0$ in F , we have

$$(53) \quad e(z)^{l-1} v_A = (l-1)! \sum_{1 \leq i_1 < \dots < i_{l-1} \leq k} e(z)_{i_1} \dots e(z)_{i_{l-1}} v_A,$$

$$e(z)_i = \underbrace{\text{Id} \otimes \dots \otimes \text{Id}}_{i-1} e(z) \otimes \text{Id} \otimes \dots \otimes \text{Id}.$$

Consequently, since $d_l \neq 0$ and $e_{\leq d} v(d) = 0$, then

$$[e(z)^{l-1}]_{\sum_{j=1}^{l-1} (l-j)d_j} v_A = (l-1)! (e_{d_1} \otimes \dots \otimes e_{d_1+\dots+d_{l-1}} \otimes \text{Id} \otimes \dots \otimes \text{Id}) v_A.$$

Lemma is proved. $\square$

Remark 4.1. Note that if $l-1 = k$ then we as a corollary obtain the proposition (3.1).

Remark 4.2. One can see that

$$(54) \quad (e_{d_1} \otimes \dots \otimes e_{d_1+\dots+d_{l-1}} \otimes \text{Id} \otimes \dots \otimes \text{Id}) v_A = v(1+d_1) \otimes v(1+d_1+d_2) \otimes \dots$$

$$\dots \otimes v(1+d_1+d_2+\dots+d_{l-1}) \otimes v(-1+d_1+\dots+d_l) \otimes \dots \otimes v(-1+d_1+\dots+d_k).$$

Denote this vector by u .

Using the fermionic realization of the modules W^A , we obtain:

Statement 4.1. a). Let $d_l \geq 2$. Denote by $A' \in \mathbb{N}^{n-2}$ such set that the associated set D' is the following: $d'_l = d_l - 2$, $d'_i = d_i$ if $i \neq l$. Then, if we will consider $W^{A'}$ as a ring $\mathbb{C}[e_0, \dots, e_{n-3}]/J^{A'}$, then there exists the isomorphism of the rings:

$$\mathbb{C}[e_2, \dots, e_{n-1}]u \cong W^{A'}, \quad e_i \mapsto e_{i-2}.$$

b). Let $d_l = 1$, $l = \min\{m : d_m \neq 0\}$, $l \neq k+1$. Let $d_{l+1} = \dots = d_{l-1} = 0$, $d_{l_1} \neq 0$. Denote by $A' \in \mathbb{N}^{n-1}$ such set that an associated set D' is of the following form:

$$d'_l = d_l - 1, d'_{l_1} = d_{l_1} - 1, d'_{l_1-l+1} = d_{l_1-l+1} + 1,$$

and for all other i $d_i = d'_i$. Then we have the isomorphism of the rings:

$$\mathbb{C}[e_1, \dots, e_{n-1}]u \cong W^{A'}, \quad e_i \mapsto e_{i-1}.$$

Remark 4.3. Note that the condition $l = \min\{m : d_m \neq 0\}$ from b) seems to be unnatural. But in the case of geneneral l the module $\mathbb{C}[e_1, \dots, e_{n-1}]u$ is not of the type $W^{A'}$ for any A' . Still, the structure of modules $\mathbb{C}[e_1, \dots, e_{n-1}]u$ in the general case is very interesting too and we hope to study it in the future.

So, there is a subspace $W^{A'}$ in W^A . Let us study the quotient. Recall that (see [1]) the dual space $(W^A)^* = (\mathbb{C}[e_0, \dots, e_{n-1}]/J^A)^*$ can be realized as a space of the polynomials $f(z_1, \dots, z_s)$, $s = 0, 1, 2, \dots$ with a following conditions:

- 1). $f(z_1, \dots, z_s)$ is symmetric.
- 2). $\deg_{z_i} f(z_1, \dots, z_s) < n$.
- 3). $f(\underbrace{z, \dots, z}_i, z_{i+1}, \dots, z_s) \div z^{\sum_{j=1}^n (i+1-a_j)_+}, i = 1, \dots, s$.

Let us prove the following proposition:

Proposition 4.1. *a). $d_l \geq 2$. Denote by $A'' \in \mathbb{N}^n$ such set that the associated set D'' is the following:*

$$d''_{l-1} = d_{l-1} + 1, d''_l = d_l - 2, d''_{l+1} = d_{l+1} + 1, d''_i = d_i \text{ otherwise.}$$

Then $W^A/W^{A'} \cong W^{A''}$.

b). $d_l = 1$, $l = \min\{m : d_m \neq 0\}$. Denote by $A'' \in \mathbb{N}^n$ such set that the associated set D'' is the following:

$$d''_{l-1} = d_{l-1} + 1, d''_l = 0, d''_{l+1} = d_{l+1} - 1, d''_{l+2} = d_{l+2} + 1, d''_i = d_i \text{ otherwise.}$$

Then $W^A/W^{A'} \cong W^{A''}$.

Proof. Note that the dual space

$$(W^A/\mathbb{C}[e_0, \dots, e_{n-1}]u)^*$$

can be realized as a space of the symmetric polynomials f in s variables ($s = 0, 1, \dots$), with a degree not more than $n - 1$ in each and satisfying the following conditions:

$$*). f(\underbrace{z, \dots, z}_i, z_{i+1}, \dots, z_s) \div z^{\sum_{j=1}^n (i+1-a_j)_+}, i = 1, \dots, s.$$

$$**). f(\underbrace{z, \dots, z}_{l-1}, z_l, \dots, z_s) \div z^{1 + \sum_{j=1}^n (l-a_j)_+}$$

(recall that $u = [e^{(n)}(z)]_{\sum_{j=1}^{l-1} (l-j)d_j} v_A$). We will prove that if the polynomial f satisfies the conditions $*)$, $**$), then it also satisfies the conditions on the function from the dual space $(W^{A''})^*$. If we also check that

$$(55) \quad \dim W^A = \dim W^{A'} + \dim W^{A''},$$

we will obtain the proof of the proposition.

Firstly, let us check that $\dim W^A = \dim W^{A'} + \dim W^{A''}$. As we have mentioned above, $\dim W^A = \prod a_i$. But then (55) is tantamount to the equality $l^2 = 1 + (l - 1)(l + 1)$ in the case *a*) and $ll_1 = l_1 - l + 1 + (l - 1)(l_1 + 1)$ in the case *b*).

Now, let us prove that if f satisfies the conditions $*)$, $**$), then $f \in (W^{A''})^*$ that is

$$(56) \quad f(\underbrace{z, \dots, z}_i, z_{i+1}, \dots, z_s) \div z^{\sum_{j=1}^i d''_j (i+1-j)_+}, i = 1, \dots, s.$$

a). $d_l \geq 2$. If $i < l - 1$ (56) is a consequence of *). If $i = l - 1$ it is a consequence from **). Let $i = l$. Condition *) gives us

$$f(\underbrace{z, \dots, z}_l, z_{l+1}, \dots, z_s) \div z^{\sum_{j=1}^l (l+1-j)d_j}.$$

But $d_{l-1}'' = d_{l-1} + 1, d_l'' = d_l - 2$. So

$$2d_{l-1}'' + d_l'' = 2d_{l-1} + d_l.$$

Thus, (56) is true for $i = l$. Now let $i > l$. Then

$$(d_{l-1} + 1)(i + 1 - (l - 1)) + (d_l - 2)(i + 1 - l) + (d_{l+1} + 1)(i + 1 - (l + 1)) = \\ = d_{l-1}(i + 1 - (l - 1)) + d_l(i + 1 - l) + d_{l+1}(i + 1 - (l + 1)).$$

So, we obtain (56) as a corollary from *). Case a) is proved.

b). $d_l = 1$. Recall that for $m < l$ we have $d_m = 0$. So for $i < l - 1$ the condition (56) is trivial.

We know from **) that

$$(57) \quad f(\underbrace{z, \dots, z}_{l-1}, z_l, \dots, z_s) \div z.$$

Using this formula, one can obtain that

$$(58) \quad f(\underbrace{z, \dots, z}_{l-1+k}, z_{l+k}, \dots, z_s) \div z^{k+1}, \quad k \geq 0.$$

This gives us (56) for $i = l - 1, \dots, l_1 - 1$. In the same time, for $i \geq l_1$ the condition (56) coincides with a condition *). Proposition is proved. $\square$

4.2. Decomposition of L^D .

Here we obtain the decomposition of L^D on the irreducible components.

Recall that we construct L^D as a limit of the inclosed spaces M^{A_s} . Note that the proposition (4.1) can be reformulated in a following way: $M^A \hookrightarrow M^{A'}$, $M^A/M^{A'} \cong M^{A''}$.

Recall that we have constructed a fermionic realization of M^A . We want to understand, how do M^A and $M^{A'}$ are embedded in $F^{\otimes k}$ with respect to each other. We know that $M^A/M^{A'} \cong M^{A''}$. In the same time, we will calculate the minimal level, in which M^A doesn't coincide with $M^{A'}$ that is the "starting" level for $M^{A''}$. It is clear that the number of this level is $\deg_q w_A - \deg_q w_{A''}$.

Let $2\alpha_i = d_{k+1} + \dots + d_{i+1}, i = 1, \dots, k$, and define $p(D)$ as a number of half-integers α_i .

Lemma 4.2. a). $d_l \geq 2, l \neq k + 1$. Then

$$\deg_q w_A - \deg_q w_{A''} = \frac{p(D'') - p(D)}{4} + \frac{d_l - 1}{2}.$$

b). $d_l = 1, l_1 = \min\{i > l : d_i \neq 0\}$. Then

$$\deg_q w_A - \deg_q w_{A''} = \frac{p(D'') - p(D)}{4} + \frac{d_{l_1}}{2}.$$

Proof. From the fermionic realization one can obtain

$$\deg_q w_A = \frac{-p(D)}{4} + \sum_{i=1}^k \alpha_i^2.$$

The rest of the proof is straightforward. $\square$

Theorem 4.1. *Let $D \in (N \cup 0)^{k+1}$, $d_1 = 0$, $d_{k+1} \leq 1$. Let $d_2 + \dots + d_{k+1} \geq 2$. Let $2 \leq l \leq k$, $d_{<l} = 0$, $l_1 = \min\{i : i > l, d_i \neq 0\}$.*

a). $d_l \geq 2$. Define D', D'' as above with an only change: if $l = k$, then $d''_{k+1} = (d_{k+1} + 1) \bmod 2$. Then

$$\text{ch}L^D = \text{ch}L^{D'} + q^{\frac{p(D'')-p(D)}{4} + \frac{d_l-1}{2}} \text{ch}L^{D''}.$$

b₁). $d_l = 1$, $l_1 \neq k+1$. Then

$$\text{ch}L^D = \text{ch}L^{D'} + q^{\frac{p(D'')-p(D)}{4} + \frac{d_{l_1}}{2}} \text{ch}L^{D''}.$$

b₂). $d_l = 1$, $l_1 = k+1$. Then

$$\text{ch}L^D = \text{ch}L^{D'}.$$

Proof. Our theorem is a consequence of the lemma (4.2). Note only that in the case b_2) the second summand vanishes, because the difference $\deg_q w_{A_s} - \deg_q w_{A'_s}$ goes to infinity while $s \rightarrow \infty$. In the other cases this difference doesn't depend on s . $\square$

Corollary 4.1. *From the equations for the characters of the integrable highest weight $\widehat{\mathfrak{sl}}_2$ -modules, we obtain the equalities for the modules itself: (all notations are from the theorem)*

a). $L^D = L^{D'} \oplus L^{D''}$. b₁). $L^D = L^{D'} \oplus L^{D''}$. b₂). $L^D = L^{D'}$.

Corollary 4.2. *We obtain an algorithm for the decomposition of the modules L^D on the irreducible components.*

Proof. Note that such D that $\sum d_i \leq 1$, correspond to the irreducible representations $L_{i,k}$. To all the rest of L^D we can apply our decomposing procedure. Note that for D, D', D'' , defined above, we have $\prod i^{d_i} > \prod i^{d'_i}$, $\prod i^{d_i} > \prod i^{d''_i}$. So, our algorithm in a finitely many steps will give us the decomposition of L^D on the sum of the irreducible $\widehat{\mathfrak{sl}}_2$ -modules. $\square$

4.3. The connection with the Verlinde algebra.

Recall the definition of the Verlinde algebra V_k , associated with a Lie algebra $\mathfrak{sl}_2$. Consider an algebra with a basis $\pi_1, \pi_2, \dots$ and with a multiplication ($i \leq j$) $\pi_i \pi_j = \pi_{j-i+1} + \dots + \pi_{i+j-1}$ (generators are multiplying as a finite-dimensional irreducible $\mathfrak{sl}_2$ -modules). By definition, $V_k = \langle \pi_1, \pi_2, \dots \rangle / (\pi_{k+1})$. Here is a list of some properties of V_k :

Statement 4.2. *The following equations are true in V_k :*

- 1). $\pi_k^2 = 1 (= \pi_1)$.
- 2). $\pi_i^2 = 1 + \pi_{i-1} \pi_{i+1}$, $i = 2, \dots, k-1$.
- 3). $\pi_i \pi_j = \pi_{j-i+1} + \pi_{i-1} \pi_{j+1}$, $i < j$, $i = 2, \dots, k-2$, $j = 3, \dots, k-1$.
- 4). $\pi_i \pi_k = \pi_{k-i+1}$, $i = 2, \dots, k-1$.

Let $D \in (\mathbb{N} \cup 0)^{k+1}$, $d_{k+1} \leq 1$. Denote

$$(59) \quad \pi_D = \pi_1^{d_1} \pi_2^{d_2} \dots \pi_{k+1}^{d_{k+1}} \in V_{k+1}.$$

We have

$$(60) \quad \pi_D = c_{1,D} \pi_1 + \dots + c_{k+1,D} \pi_{k+1}.$$

From the corollaries 4.1, 4.2 and statement 4.2 we obtain the following proposition:

Proposition 4.2. *There is a decomposition*

$$L^D = M_1 \otimes L_{0,k} \oplus \dots \oplus M_{k+1} \otimes L_{k,k},$$

where $\dim M_i = c_{i,D}$ and M_i are graded spaces. The character of M_i can be obtained by the procedure, described above.

5. COMBINATORIAL CALCULATIONS

In this section we will obtain the formulas for the character of L^D . As a consequence we will obtain the formula from [6] for the characters of the irreducible representations $L_{i,k}$. Using this formulas, we will reprove an equation a) from the corollary (4.1).

5.1. The character formula for L^D .

One can see that the spaces L^D are bigraded by the operators h_0 and d (for $x \in \mathfrak{sl}_2$ $[d, x_i] = ix_i$). So, one can introduce a character as a trace $\text{Tr} q^d z^{h_0}$. In the same time, it is clear that for all D the eigenvalues of h_0 on L^D are of the same parity. So, we will define the character in the following way: $\text{ch} L^D = \text{Tr} q^d z^{\frac{h_0}{2}}$. In this definition, all the powers of z are integer or half-integer. (One can see that they are integer, if $\sum_{i=2}^{k+1} (i-1)d_i$ is even, and half-integer, if it is odd).

Recall that in [1] we obtain the following formula for the character of W^A ($A = (a_1 \leq \dots \leq a_n)$, $a_n = k+1$, $d_j = \#\{i : a_i = j\}$, $d_1 = 0$):

$$(61) \quad \text{ch}(W^A, q, qz) = \sum_{j_k=0}^{d_{k+1}} \sum_{j_{k-1}=0}^{d_k+j_k} \dots \sum_{j_1=0}^{d_2+j_2} z^{\sum_{l=1}^k j_l} q^{\sum_{l=1}^k j_l(d_2+\dots+d_l+j_l)} \times \\ \times \begin{bmatrix} d_{k+1} \\ j_k \end{bmatrix}_q \begin{bmatrix} d_k+j_k \\ j_{k-1} \end{bmatrix}_q \dots \begin{bmatrix} d_2+j_2 \\ j_1 \end{bmatrix}_q.$$

Let us now obtain the character of M^A (again, powers of z will be integer or half-integer). Denote

$$2\alpha_i = d_{k+1} + \dots + d_{i+1} \quad i = 1, \dots, k.$$

In this notations $p(D) = \#\{i : \alpha_i \in (\mathbb{Z} + \frac{1}{2})\}$. Recall that

$$W^A = \mathbb{C}[e_0, \dots, e_{n-1}]v_A, \quad M^A = \mathbb{C}[e_0, \dots, e_{-n+1}]w_A.$$

Thus, to obtain M^A from W^A one must move v_A to w_A and then "turn" W^A , to get e_{1-i} from e_{n-i} . One can check that while moving $v_A \rightsquigarrow w_A$ the z -degree will decrease on $\sum \alpha_i$, and the q -degree will increase on $\sum \alpha_i^2 - \frac{p(D)}{4}$ (note that in the formula (61) we regard the q -degree of v_A to be equal to zero, so $\sum \alpha_i^2 - \frac{p(D)}{4}$ is just a q -degree of w_A in $F^{\otimes k}$). Thus, to obtain from (61) the formula for the character of M^A , one must multiply (61) on the corresponding factor and replace z by $\frac{z}{q^n}$, what is corresponding to the "turn" of W^A (we must divide on q^n , because the formula for the character of W^A depends on the variables (q, zq)). We obtain the

following expression (substituting $n = \sum d_i$):

$$(62) \quad \text{ch}(M^A, q, z) = z^{-\sum_{i=1}^k \alpha_i} q^{\sum_{i=1}^k \alpha_i^2} q^{-p(D)/4} \times \\ \times \sum_{j_k=0}^{d_{k+1}} \sum_{j_{k-1}=0}^{d_k+j_k} \cdots \sum_{j_1=0}^{d_2+j_2} \left(\frac{z}{q^{\sum_{i=2}^{k+1} d_i}} \right)^{\sum_{l=1}^k j_l} q^{\sum_{l=1}^k j_l (d_2 + \cdots + d_l + j_l)} \times \\ \times \begin{bmatrix} d_{k+1} \\ j_k \end{bmatrix}_q \begin{bmatrix} d_k + j_k \\ j_{k-1} \end{bmatrix}_q \cdots \begin{bmatrix} d_2 + j_2 \\ j_1 \end{bmatrix}_q.$$

After a simple rearranging, we obtain:

$$(63) \quad \text{ch}(M^A, q, z) = q^{-p(D)/4} \times \\ \times \sum_{j_k=0}^{d_{k+1}} \sum_{j_{k-1}=0}^{d_k+j_k} \cdots \sum_{j_1=0}^{d_2+j_2} z^{\sum_{l=1}^k (j_l - \alpha_l)} q^{\sum_{l=1}^k (j_l - \alpha_l)^2} \times \\ \times \begin{bmatrix} d_{k+1} \\ j_k \end{bmatrix}_q \begin{bmatrix} d_k + j_k \\ j_{k-1} \end{bmatrix}_q \cdots \begin{bmatrix} d_2 + j_2 \\ j_1 \end{bmatrix}_q.$$

Make a change $i_l = j_l - \alpha_l$:

$$(64) \quad \text{ch}(M^A, q, z) = q^{-p(D)/4} \times \\ \times \sum_{\substack{i_k = -\alpha_k \\ i_k + \alpha_k \in \mathbb{Z}}}^{d_{k+1}/2} \sum_{\substack{i_{k-1} = -\alpha_{k-1} \\ i_{k-1} + \alpha_{k-1} \in \mathbb{Z}}}^{d_k/2 + i_k} \cdots \sum_{\substack{i_1 = -\alpha_1 \\ i_1 + \alpha_1 \in \mathbb{Z}}}^{d_2/2 + i_2} z^{\sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2} \times \\ \times \begin{bmatrix} d_{k+1}/2 + \alpha_k \\ i_k + \alpha_k \end{bmatrix}_q \begin{bmatrix} d_k/2 + i_k + \alpha_{k-1} \\ i_{k-1} + \alpha_{k-1} \end{bmatrix}_q \cdots \begin{bmatrix} d_2/2 + i_2 + \alpha_1 \\ i_1 + \alpha_1 \end{bmatrix}_q.$$

Rewrite the binomial coefficients:

$$(65) \quad \text{ch}(M^A, q, z) = q^{-p(D)/4} \sum_{\substack{i_l \geq -\alpha_l, \ i_l + \alpha_l \in \mathbb{Z} \\ i_k \leq \frac{d_{k+1}}{2}, \ i_{l+1} \geq i_l - \frac{d_{l+1}}{2}}} z^{\sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2} \times \\ \times \frac{1}{(i_k + \frac{d_{k+1}}{2})_q! \prod_{l=1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!} \times \\ \times \frac{(d_{k+1})_q!}{(\frac{d_{k+1}}{2} + i_k)_q!} \prod_{l=1}^{k-1} \frac{(i_{l+1} + \frac{d_{l+1}}{2} + \alpha_l)_q!}{(i_l + \alpha_l)_q!}.$$

We are interesting in the limit of (65) while $d_{k+1} \rightarrow \infty$ (d_{k+1} must preserve its parity). Denote

$$(\infty)_q = \prod_{i=1}^{\infty} (1 - q^i).$$

(In our formulas the symbol $(\infty)_q$ for the $\prod_{i=1}^{\infty} (1 - q^i)$ seems to be more natural, than the more common one $(q)_{\infty}$). In order to compute the limit, note that we can

consider such i_l that $-d_{k+1}/4 < i_l < d_{k+1}/4$ (otherwise the q -degree will overreach $d_{k+1}^2/16$). Then the degree in q of the polynomial

$$\frac{(i_{l+1} + \frac{d_{l+1}}{2} + \alpha_l)_q!}{(i_l + \alpha_l)_q!} - 1$$

is greater than $\frac{d_{k+1}}{4}$. So

$$\frac{(d_{k+1})_q!}{(\frac{d_{k+1}}{2} + i_k)_q!} \prod_{l=1}^{k-1} \frac{(i_{l+1} + \frac{d_{l+1}}{2} + \alpha_l)_q!}{(i_l + \alpha_l)_q!} \longrightarrow 1 \text{ while } d_{k+1} \rightarrow \infty.$$

In the same time

$$\frac{1}{(i_k + \frac{d_{k+1}}{2})_q!} \longrightarrow \frac{1}{(\infty)_q}.$$

We have proved the following theorem:

Theorem 5.1. *Let $D \in (\mathbb{N} \cup 0)^{k+1}$, $d_{k+1} \leq 1$. Denote $2\alpha_i = d_{k+1} + \dots + d_{i+1}$, $i = 1, \dots, k$. Then*

$$\text{ch}L^D = \frac{q^{-p(D)/4}}{(\infty)_q} \sum_{\substack{i_1, \dots, i_k \\ i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l=2, \dots, k}} \frac{z^{\sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2}}{\prod_{l=1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!}.$$

Corollary 5.1. *We obtain the formula from [6]:*

$$\text{ch}L_{j,k} = \frac{1}{(\infty)_q} \sum_{\substack{i_1, \dots, i_k \in \mathbb{Z} \\ i_1 \leq \dots \leq i_k}} \frac{z^{\frac{j}{2} + \sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2 + \sum_{l=1}^j i_l}}{\prod_{l=1}^{k-1} (i_{l+1} - i_l)_q!}.$$

5.2. Combinatorial recurrent formula for L^D . Introduce the following notations:

$$(66) \quad D = (\dots d_{s-1}(d_s + 2)d_{s+1} \dots), D' = (\dots d_{s-1}d_sd_{s+1} \dots), \\ D'' = (\dots (d_{s-1} + 1)d_s(d_{s+1} + 1) \dots).$$

Proposition 5.1. $\text{ch}L^D = \text{ch}L^{D'} + q^{\frac{p(D'') - p(D)}{4} + \frac{d_s + 1}{2}} \text{ch}L^{D''}$.

Proof. Note that $p(D) = p(D')$. We have:

$$(67) \quad \text{ch}L^D - \text{ch}L^{D'} = \frac{q^{-p(D)/4}}{(\infty)_q} \times \\ \times \left(\sum_{\substack{i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_s \geq i_{s-1} - \frac{d_s + 2}{2} \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l \neq s}} \frac{z^{\sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2}}{(i_s - i_{s-1} + \frac{d_s + 2}{2})_q! \prod_{l=1, l \neq s-1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!} - \right. \\ \left. - \sum_{\substack{i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l=2, \dots, k}} \frac{z^{\sum_{l=1}^k i_l} q^{\sum_{l=1}^k i_l^2}}{\prod_{l=1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!} \right).$$

Divide the summation region of the minuend into the two parts: the first one with a parameters with $i_{s+1} = i_s - \frac{d_s+2}{2}$, and the second one with $i_{s+1} \geq i_s - \frac{d_s}{2}$. Then the conditions on the parameters in the second part will coincide with the conditions on the parameters in subtrahend. Thus the previous expression equals

$$(68) \quad \frac{q^{-p(D)/4}}{(\infty)_q} \left(\sum_{\substack{i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_s = i_{s-1} - \frac{d_s+2}{2} \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l \neq s}} \sum_{l=1}^k i_l \sum_{l=1}^k i_l^2 \frac{1}{\prod_{l=1, l \neq s-1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!} + \right. \\ \left. + \sum_{\substack{i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l=2, \dots, k}} \sum_{l=1}^k i_l \sum_{l=1}^k i_l^2 \frac{q^{i_s - i_{s-1} + \frac{d_s+2}{2}}}{(i_s - i_{s-1} + \frac{d_s+2}{2})_q! \prod_{l=1, l \neq s-1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!} \right).$$

Add the first summand to the second:

$$(69) \quad \frac{q^{-p(D)/4}}{(\infty)_q} \sum_{\substack{i_l + \alpha_l \in \mathbb{Z}, l=1, \dots, k \\ i_s \geq i_{s-1} - \frac{d_s+2}{2} \\ i_l \geq i_{l-1} - \frac{d_l}{2}, l \neq s}} \sum_{l=1}^k i_l \sum_{l=1}^k i_l^2 \frac{q^{i_s - i_{s-1} + \frac{d_s+2}{2}}}{(i_s - i_{s-1} + \frac{d_s+2}{2})_q! \prod_{l=1, l \neq s-1}^{k-1} (i_{l+1} - i_l + \frac{d_{l+1}}{2})_q!}.$$

Let us change the parameters $i_s := i_s + 1/2$, $i_{s-1} := i_{s-1} - 1/2$. Then we obtain:

$$\text{ch} L^D - \text{ch} L^{D'} = q^{\frac{p(D'') - p(D) + 2d_s + 2}{4}} \text{ch} L^{D''}.$$

Proposition is proved. $\square$

Corollary 5.2. $L^D = L^{D'} \oplus L^{D''}$.

REFERENCES

- [1] B. Feigin, E. Feigin, Q-characters of the tensor product in $\mathfrak{sl}_2$ -case, q-alg 0201111.
- [2] B. Feigin, S. Loktev, On generalized Kostka polynomials and quantum Verlinde rule, q-alg 9812093.
- [3] B. Feigin, S. Loktev, On finitization of Gordon identities, *Funct. Anal. Appl.* **35**, 53-61 (2001).
- [4] Anne Schilling, S. Ole Warnaar, Supernomial coefficients, polynomial identities and q-series, q-alg 9701007
- [5] S. O. Warnaar, The Andrews-Gordon identities and q-multinomial coefficients, q-alg 9601012
- [6] B. Feigin, A. Stoyanovski, Quasi-particles models for the representations of Lie algebras and geometry of flag manifold, hep-th/9308079, RIMS 942; Functional models for the representations of current algebras and the semi-infinite Schubert cells, *Funct. Anal. Appl.* **28** (1994), 55-72.
- [7] B. Feigin, T. Miwa, Extended vertex operator algebras and monomial bases, q-alg 9901067.
- [8] B. Feigin, R. Kedem, S. Loktev, T. Miwa, E. Mukhin, Combinatorics of the $\widehat{\mathfrak{sl}_2}$ spaces of coinvariants, q-alg 9908003.
- [9] B. Feigin, R. Kedem, S. Loktev, T. Miwa, E. Mukhin, Combinatorics of the $\widehat{\mathfrak{sl}_2}$ spaces of coinvariants II, q-alg 0009198.
- [10] A. N. Kirillov, M. Shimozono, A generalization of Kostka-Foulkes polynomials, q-alg 9803062.
- [11] A. Schilling, S. Ole Warnaar, Inhomogeneous lattice paths, generalized Kostka polynomials and A_{n-1} supernomials, q-alg 9802111.

B.F.: LANDAU INSTITUTE FOR THEORETICAL PHYSICS, RUSSIA, CHERNOGOLOVKA, 142432
E-mail address: feigin@mccme.ru

E.F.: INDEPENDENT UNIVERSITY OF MOSCOW, RUSSIA, MOSCOW, BOL'SHOI VLAS'EVSKII PER., 7.
E-mail address: evgfeig@mccme.ru